

Antimicrobial Activities from the Essential Oil from Leaves of Eucalyptus Globulus Against the Microbes Isolated from Contaminated Soil, Water & Milk

Pragnya paramita Acharya², Ajay Kumar Sahu¹, Barsha Rani Kar²

¹Department of Microbiology, Bangalore University, Bangalore-560043, India

²Department of Biotechnology, Academy of management & Information Technology Khordha- 52057, Odisha, India

Abstract - Eucalyptus species are well known as medicinal plants because their biological and pharmacological properties. In the international pharmacopeia, the most important and represented species, used anaesthetic, anodyne, antiseptic, astringent, deodorant, diaphoretic, disinfectant, expectorant, febrifuge, fumigant, haemostat, inhalant, insect repellent, preventive, rubefacient, sedative yet abscess, arthritis, asthma, bolus, bronchitis, burns, cancer, diabetes, diarrhea, diphtheria, dysentery, encephalitis, enteritis, erysipelas, fever, flu, inflammation, laryngalgia, laryngitis, leprosy, malaria, mastitis, trachalgio, worms and wounds, some time their demand is also high in the soap and cosmetic industries.

Keywords:- Antimicrobial, Essential oils, Eucalyptus leaves, Isolated microbes, Cancer activities.

INTRODUCTION

Eucalyptus globulus belongs to the Myrtaceae family, native to Australia (Mota et al., 2015). It is tall evergreen tree, it was introduced in Algeria in 1854 by Rame (Bachir Raho & Benali, 1999). This oil has been used especially to manufacture pharmaceutical inhalants, nasal discharge stimulants, oral care products, or even with the function of flavor and aroma to medicines. However, recent evidence points to possible effects associated with healing, anti-inflammatory and antimicrobial action (Mota et al., 2015). Eucalyptus species are well known as medicinal plant because of their biological and pharmacological properties. In the international pharmacopeia, the most important and represented species, used anaesthetic, anodyne, antiseptic, astringent, deodorant, diaphoretic, disinfectant, expectorant, febrifuge, fumigant, hemostat, inhalant, insect repellent, preventive, rubefacient, sedative yet abscess, arthritic, asthma, boils, bronchitis, burns, cancer, diabetes, diarrhea, diphtheria, diphtheria, dysentery, encephalitis, enteritis, erysipelas, fever, flu, inflammation, laryngalgia, laryngitis, leprosy, malaria, mastitis, miasma, phasms, trachalgio, worms, and wounds, some time their demand is also high in the soap and cosmetic industries (Bachir Raho, 2012).

Leaf extracts of Eucalyptus have been recognized as natural food preservatives (Mori, 2004). All aforementioned products are widely cultivated in Brazil and evaluating the potential of these substances can result in future investigations hypotheses. For example, the development of antiseptic or disinfectant products to address the emerging risk of reduction in sensitivity of the microorganisms to the germicidal agents. Another example is to stimulate discoveries related to the prevention and treatment of bio-films that slow or prevent the healing of wounds (especially in chronic wounds), generating high costs for treatment and associated morbidity and mortality. Therefore, the present study sought to evaluate the in vitro antimicrobial activity of the essential oil of Eucalyptus globulus, of xylitol against the following microorganisms: Pseudomonas aureginosa; Salmonella sp.; Staphylococcus aureus; Proteus vulgaris; Escherichia coli and Candida albicans. (Valéria de Siqueira Mota et al.). Reports have shown that about 50,000 people die in the world per day due to infectious diseases mainly caused by Candida albicans, Escherichia coli, Pseudomonas aeruginosa and Staphylococcus aureus. (Muhammad Rifaqat Ammer et al.). Eucalyptus (EO) has antibacterial, antiviral, antifungal components and a long history of use against the effect of colds, influenza, other respiratory infection, rhinitis, and sinusitis (Angela E. Sadlon et al.). Escherichia coli, originally called "Bacterium coli" was first isolated from the faces of a child in 1885 by the Austrian pediatrician Theodor Escherich (Escherich, 1885). Escherichia coli is a common inhabitant of the gastrointestinal tract of humans and animals. There are E. coli strains that are harmless in the intestinal tract. The pathogenic E. coli are divided into those strains causing disease inside the intestinal tract and others capable of infection at extra-intestinal sites (Kaper et al., 2004). Escherichia coli can be found secondarily in soil and water as the result of fecal contamination, E. coli may freely replicate in tropical fresh water (Bermudez and Hazen, 1988). Classically, its detection has been used as an indicator of poor water quality

(Rodney A. Welch). The ability to build a phylogeny for some of the E. coli pathotypes is complicated because the horizontally acquired islands do not always share chromosomal location and genetic content (Welch et al., 2002). Escherichia coli are Gram-negative, non-spore-forming bacilli. They are approximately 0.5 µm in diameter and 1.0–3.0 µm in length. Within the periplasm is a single layer of peptidoglycan, peptidoglycan has a typical subunit structure where the N-acetylmuramic acid is linked by an amide bond to a peptide consisting of L-alanine, D-glutamic acid, meso-diaminopimelic acid and finally D-alanine. Escherichia coli are commonly motile in liquid by means of peritrichous flagella. Swarming behavior and differentiation into hyperflagellated and elongated bacilli typical of that seen with the Proteus species can be observed on some solid media (Harshey 1994). Most E. coli strains are capable of growing over a wide range in temperature (approximately 15–48°C). The growth rate is maximal in the narrow range of 37–42° (C Ingraham and Marr, 1987). Escherichia coli can grow within a pH range of approximately 5.5–8.0 with best growth occurring at neutrality. Some diarrheagenic E. coli strains have the ability to tolerate exposure to pH 2.0. Such an acid shock mimics transit through the stomach and induces expression of sets of genes involved in survival and pathogenesis (Waterman and Small, 1996). Escherichia coli provide many necessary vitamins including Vitamin K and B-complex vitamins. We have billions of E. coli bacteria in our bodies, making things we need, helping digest our food and maintaining our health. Although most E. coli are harmless and are a needed bacterium for health, there are some strains of E. coli bacteria that can be very harmful to our health. A rare strain of E. coli that you may have seen in the news can cause potentially dangerous outbreaks and illness. This strain is E. coli O157:H7. This E. coli can produce a toxin called Shiga-like toxin (SLT). The harmful strain of E. coli bacteria can cause abdominal cramping, diarrhoea and occasionally vomiting. Usually little or no fever is present. Dehydration, even in mild cases of diarrhoea, can easily occur. Normally the illness resolves in 5 to 10 days. In 5%-10% of cases, hemolytic uremic syndrome (HUS), which is characterized by kidney failure and loss of red blood cells, can occur. In severe cases of the disease, 2%-7% may have permanent kidney damage. Dehydration is particularly dangerous to small children who are too small to tolerate much blood and fluid loss. The presence of these bacteria can also be very dangerous to the elderly population or persons who are already ill (Centers for Disease Control and Prevention, National Center for Infectious Diseases, Division of Bacterial and Mycotic Diseases. January 27, 2004).

Staphylococcus aureus is Gram-positive, Cocci in grape-like clusters (Sarkis K. Mazmanian, 7 FEBRUARY 2003). Staphylococcus aureus is one of the most common causes of community- and health care-associated infections, Staphylococcus aureus is one of the most successful human pathogens, little is known about its effects on the U.S. population as a whole. We analyzed newly available data from the 2001–2002 National Health and Nutrition Examination Survey (NHANES). (Philip L. Graham III, 7 March 2006). Staphylococcus aureus accounts for about 13% of all nosocomial blood infections, and is the second most common cause of these infections after coagulase negative staphylococci. 1 S. aureus bacteraemia extends length of hospital stay and increases antibiotic use, costs, and mortality. 1 Nasal carriage is a risk factor for acquiring nosocomial infection. 2 It has been shown that 80% of nosocomial S. aureus bacteraemia episodes in carriers of the bacteria are attributable to an endogenous source. (Heiman F L Wertheim, et al Vol 364 August 21, 2004). The gram positive bacterium S. aureus is mainly responsible for postoperative wound infection. Expression of virulence determinants in S. aureus. It is known for its capacity to cause a broad range of important infections in humans. Such capacity is related to the expression of an array of factors that participate in pathogenesis of infection, allowing this bacterium to adhere to surfaces/tissues, avoid or invade the immune system, and cause harmful toxic effects to the host. These factors are known as virulence determinants (Table 1), and can be divided into cell-surface-associated (adherence) and secreted (exotoxins) factors. Table 1 Virulence factors involved in the pathogenesis of Staphylococcus aureus and respective putative function. (Ana Rita Costa. 2013).

Table-1

VIRULENCE FACTOR	PUTATIVE FUNCTION
CELL SURFACE FACTORS Microbial surface components recognizing adhesive matrix molecules (MSCRAMMs) Staphylococcal protein A (SpA)	Bind to IgG, interfering with opsinization and phagocytosis
Fibronectin-binding proteins (FnbpA and FnbpB)	Attachment to fibronectin and plasma clot
Collagen-binding protein	Adherence to collagenous tissues and Cartilage
Clumping factor proteins (ClfA and ClfB)	Mediate clumping and adherence to fibrinogen in the presence of fibronectin
SECRETED FACTORS Superantigens Staphylococcal enterotoxins (SEA, B, C, D, E, G and Q) Toxic shock syndrome toxin-1 (TSST-1) Cytolytic toxins Cytolysins α-hemolysin β-hemolysin γ-hemolysin Leukocidin family Leukocidins E/D and M/F-PV Various exoenzymes Lipases Nucleases	Massive activation of T cells and antibody presenting cells Massive activation of T cells and antibody presenting cells Induce lysis on a wide spectrum of cells, mainly platelets and monocytes Hydrolysis of sphingomyelin of the plasmatic membrane of monocytes, erythrocytes, neutrophils and lymphocytes; make cells susceptible to other lytic agents Induce lysis on erythrocytes and Leukocytes Induce lysis on leukocytes Induce lysis on leukocytes Inactivate fatty acid Cleave nucleic acids Inactivate neutrophil activity; activate T cells (only ETA and ETB)

Proteases	
Serine (e.g. exfoliative toxins ETA and ETB)	Block neutrophil activation and Chemotaxis
Cysteine (e.g. staphopain)	Inactivate antimicrobial peptides
	Degrade hyaluronic acid
Aureolysin	Activate plasminogen; inactivate antimicrobial peptides
Hyaluronidase	
Staphylokinase (SAK)	
Miscellaneous proteins	Inhibit complement activation
Staphylococcal complement inhibitor (SCIN)	Inhibit complement activation
Extracellular fibrinogen binding protein (Efb)	Inhibit chemotaxis and activation of Neutrophils
Chemotaxis inhibitory protein of <i>S. aureus</i> (CHIPS)	Inhibit chemotaxis of neutrophils
Formyl peptide receptor-like 1 inhibitory protein (FLIPr)	
Extracellular adherence protein (Eap)	Inhibit neutrophil migration

2. MATERIAL

The Eucalyptus leaves were collected from rural area Debagarh district and from urban area Bhubaneswara. It was crushed manually with wooden arrangement and made into small pieces. The bacterial samples were collected from 3 different places (i) drain water collected from drain of AMIT, (ii) soil collected from near the dustbin, (iii) the unpasteurized raw milk was collected from the cattle house of Khordha.



11 2 (fig- Eucalyptus leaves were collected from 1.rural area and 2.urban area)

3.METHOD

Extraction of oil was carried out by Suxhlet apparatus and it was working on steam distillation process. 20gm pieces of leaves were taken and 200ml distilled water were used and working temperature is maintained at 100 degree Celsius.



(fig-Suxhlet apparatus)

and distilled for 1 hour. Once the distillation started the sample started boiling and vapour is formed, the vapour was cooled down with the help of condensed. The condensed material was collected on the other side of setup. The collected material is a mixture of oil and plant extract. After the water was separated by the rotary evaporator. The Eucalyptus oil was purified which was used as antimicrobial material.

Confirmation test of presence of oil in the plant extract by grease paper test. Two pieces of filter paper were taken, then a drop of rural area oil on first paper and urban area oil on another paper piece. The paper did not absorb the thick plant extract which proves the presence of oil.

Confirmation test for presence of oil in the plant extract by chemical method, a few drops of Phenolphthalein were taken in a test tube then added two drops of alkali solution (NaOH) and its colour was converted into pink colour.


Then the plant extracted oil was added on the pink colour solution and its colour converted into white which indicated the presence of fatty acid. The extracted oil was kept in 4 degree Celsius freezer. Then 2.8gm of nutrient agar was dissolved in 200 ml of distilled water and 2.6gm of nutrient broth was dissolved in another 200ml of distilled water and both solution were closed with cotton block and to autoclave at 121 degree Celsius. The will sterilized nutrient agar was transformed into 3 plates and allow incubation for 24 hours. Then after 24 hours nutrient broth transformed in to 3 test tubes then the microbes culture were transformed into the nutrient broth and allow them to incubation for 24 hours.

For gram staining, after 24 hours the microbes culture inoculate in a sterilized slide was heat fixed and an staining try, then the smear was wetted with crystal violet gently and allowed to stand for 1 minute, then the slide was slightly tilted and gently rinsed with tap water using a wish bottle, then the smear was gently flooded with Gram's iodine and again allowed to stand for one minute, the slide was slightly tilted and gently rinsed with tap water, the smear appear as a purple circle on the slide, then 95% ethyl alcohol was used to decolourize, the slide was slightly tilted until the alcohol run almost clear. the slid was immediately rinsed, then it was gently flooded with safranine to stand for 45 second. Then slid was gently rinsed with tap water, then slid was dried with bibulous paper, then the smear was viewed using a light microscope under oil immersion.

For the antimicrobial screening, numbers of small rounded filter paper were taken and wetted with plant extracted oil and inoculated in to the freshly made nutrient agar plates in which the respective microbes were inoculated for grown.

4. RESULT

Result of oil extraction



12(fig-plant extraction from 1.rural 2.urban)

Result of grease paper test



(fig- grease paper test) . The paper did not absorb the thick plant extraction which prove the presence of grease.

Result of plant extract by chemical method



(fig-before addition of plant extraction oil)



(fig- after addition of plant extraction oil)

We found that the plant extraction is a fatty aide.

Result of microorganisms grow



(fig- microorganisms grow in nutrient agar)

Result of gram staining

No. of slide sample taken	Colour observation	Shape	Character
1.Drain water	Purple	Round	Gram +ve
2. Drain water	Purple	Round	Gram +ve
3.soil	Pink	Rod	Gram -ve
4. soil	Pink	Rod	Gram -ve
5.milk	Purple	Round	Gram +ve
6. milk	Purple	Round	Gram +ve

Result of antimicrobial screening



(fig- antimicrobial screening)

Rain water

sample	Zone in cm
Rural area	1
Urban area	1.5
Milk sample	
Rural area	0.8
Urban area	1.8

Soil sample

Rural area	1.8
Urban area	1

From the experiment we found that plant extraction is acted as antimicrobial oil.

5.CONCLUSIONS

The addition of essential oil leaves in broth culture inoculated with *S. aureus* and *E. coli* inhibited the growth of these organisms. The rate of inhibition was greater, on gram negative bacteria (*E. coli*) than that observed on gram positive bacterium (*S. aureus*). In most cases the size of inoculum and the concentration of essential oil leaves affected the growth/survival of the organisms. The growths of tested bacteria in high concentrations of essential oil leaves were highly inhibited, where it was considered that these organisms were sensitive to the oil. Some authors have reported that gram-negative microorganisms are slightly more sensitive to essential oils when compared to gram-positive. The gram-positive and gram-negative microorganisms differ in several aspects other than with respect to the structure of their cellular walls, mainly with regard to the presence of lipoproteins and lipopolysaccharides in gram-negative bacteria that form a barrier to hydrophobic compounds. Some researchers reported that there is a relationship between the chemical structures of the most abundant in the tested essential oil and the antimicrobial activity. The antibacterial activity of Eucalyptus extracts has been due to the components such as 1,8-cineole, citronellal, citronellol, citronellyl acetate, p-cymene, eucamalol,

limonene, linalool, β - pinene, γ - terpinene, α - terpinol, alloocimene and aromadendrene. The essential oils from the leaf of *E. globulus* showed varying degrees of antibacterial activity against two clinical isolates. From the above experiment it can be inferred that extract suggest significant growth inhibiting effect on Gram-positive (*E. coli*) and Gram-negative bacteria (*S. aureus*). The efficacy of leaf oil of *E. globulus* against these microorganisms may provide a scientific ground for the application of the herb in the prevention and treatment of bacterial infections caused by various pathogenic bacteria such as

Staphylococcus aureus and Escherichia coli, which have developed resistance to antibiotics. The incorporation of this oil into the drug formulations is also recommended.

REFERENCES

1. Valeria de Siqueira Mota, Ruth Natalia Teresa Turrini & Vanessa de Brito Poveda, Antimicrobial activity of Eucalyptus globulus oil, xylitol and papain: a pilot study, Journal of school of nursing USP, 49, 215-219, 2015.
2. Journal of Radiation Research and Applied Sciences 9(2016) Agela E.Sadlon:etal 2010
3. Bachir Raho G & Benali M 2012 by the Asian Pacific Journal of Tropical Biomedicine
4. (Escherich, Role of respiratory syncytial virus in lower respiratory tract infections in a pediatric hospital in Mexico city. Bol Med Hosp Infant Mex 1885).
5. Kaper et al., Influenza related hospitalizations among children in Hong Kong. N Engl J Med 2002;347:2097-103
6. (RODNEY A. WELCH An epidemiological study of RSV infection in the Gambia. Bull WHO 2002;80: 562-8.
7. (Welch et al., European Antimicrobial Resistance Surveillance System Participants. European Antimicrobial Resistance Surveillance System (EARSS): susceptibility testing of invasive Staphylococcus aureus. Euro Surveill. ;5:34-6. 2002).
8. (Harshey, Drug resistance among clinical isolates of frequently encountered bacterial species in central Europe during 1975-1995. Infection. 1999;27(Suppl 2):S2-8.
9. (Waterman and Small, Staphylococcus aureus (MRSA): a community-based prevalence survey. Epidemiol Infect. 2001;126:351-6. t
10. (The Genus Escherichia, RODNEY A. WELCH Prokaryotes (2006) 6:60-71
11. (Centers for Disease Control and Prevention, National Center for Infectious Diseases, Division of Bacterial and Mycotic Diseases. January 27, 2004)
12. Passage of Heme-Iron Across the Envelope of Staphylococcus aureus (Sarkis K. Mazmanian, Eric P. Skaar, Andrew H. Gaspar, Munir Humayun, Piotr Gornicki, Joanna Jelenska, Andrzej Joachmiak, Dominique M. Missiakas, Olaf Schneewind. 7 FEBRUARY 2003 VOL 299 SCIENCE www.sciencemag.org)
13. U.S. Population-Based Survey of Staphylococcus aureus Colonization Philip L. Graham III, MD, MSc; Susan X. Lin, DrPH; and Elaine L. Larson, PhD, 7 March 2006 Annals of Internal Medicine Volume 144.
14. Risk and outcome of nosocomial Staphylococcus aureus bacteraemia in nasal carriers versus non-carriers (Margreet C Vos, Alewijn Ott, Alex van Belkum, Andreas Voss, Jan A J W Kluytmans, Peter H J van Keulen, Christina MJE Vandenbroucke-Grauls, Marlene H M Meester, HenAri Verbrugh. Vol 364, 21st August 2004 .
15. Staphylococcus aureus virulence factors and disease . Ana Rita Costa, Deivid W. F. Batistão, Rosineide M. Ribas, Rosineide M. Ribas, Ana Margarida Sousa, Olivia Pereira and Claudia M. Botelho., FORMATEX 2013.
16. Antibacterial activity of the essential oils from the leaves of Eucalyptus globulus against Escherichia coli and Staphylococcus aureus, (Bachir Raho G and Benali M) doi:10.1016/S2221-1691(12)60220-2 2012 by the Asian Pacific Journal of Tropical Biomedicine. All rights reserved.